

Letter to the editor:
REPRODUCTIVE TOXICITY OF BORON

Florian Seidel

Leibniz Research Centre for Working Environment and Human Factors, Ardeystr. 67,
44139 Dortmund, Germany, E-mail: seidelf@ifado.de

<http://dx.doi.org/10.17179/excli2020-2757>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>).

Dear Editor,

Recently, Bolt and colleagues published a review article on possible effects of boron on human reproduction (Bolt et al., 2020). The metalloid boron is required as an additive of fiber-glass, structural material and ceramics. Human exposure typically occurs in mining. Since more than a decade, a discussion is ongoing concerning categorization of inorganic boron compounds as reproductive toxins (Moore, 1997; Hasegawa et al., 2013; ECHA, 2010; SCCS, 2010; ATSDR, 2010). The authors summarized key findings relevant for this discussion (Bolt et al., 2020):

- Boron causes testicular toxicity in rats, mice and dogs.
- The NOAEL in rats was 17.5 mg boron/kg body weight per day; at higher concentrations reduced epididymal sperm counts were observed.
- The NOAEL of developmental toxicity in rats was 9.6 mg boron/kg body weight per day; at higher doses fetal skeletal effects were observed.
- The boron blood concentrations in rats corresponding to the NOAEL of 9.6 mg/kg/day was in the range of 1,200 ng boron/g.
- Even for extreme occupational exposure groups in China and Turkey boron blood concentrations were clearly below 1,200 ng/g.

Evaluation of reproductive and developmental toxicity remains a major challenge (Sachinidis et al., 2019; Leist et al., 2017; Shinde et al., 2015, 2016, 2017). Currently, several research projects aim at the development of *in vitro* tests of developmental toxicity (Waldmann et al., 2017; Rempel et al., 2015; Balmer et al., 2014). For this purpose, human stem cells have been exposed to test compounds to study if their differentiation is compromised (Weng et al., 2014; Waldmann et al., 2014; Krug et al., 2013). However, although mechanisms can be evaluated in these *in vitro* systems, it still remains challenging to extrapolate to the *in vivo* situation and derive e.g. NOAELs. Therefore, the evaluation of possible human developmental effects of boron still depends on animal experiments and as far as available on epidemiological studies. The systematic review of Bolt and colleagues suggests that occupational exposure of humans to boron results in boron blood concentrations that are clearly below levels that cause adverse effects in animal experiments.

Conflict of interest

The author declares no conflict of interest.

REFERENCES

- ATSDR (Agency for Toxic Substances and Disease Registry). Toxicological profile for boron. Atlanta, GA, USA, Nov 2010. <https://www.atsdr.cdc.gov/ToxProfiles/TP.asp?id=453&tid=80>. Accessed 7 Jan 2020.
- Balmer NV, Klima S, Rempel E, Ivanova VN, Kolde R, Weng MK, et al. From transient transcriptome responses to disturbed neurodevelopment: role of histone acetylation and methylation as epigenetic switch between reversible and irreversible drug effects. *Arch Toxicol*. 2014;88:1451-68. doi: 10.1007/s00204-014-1279-6.
- Bolt HM, Başaran N, Duydu Y. Effects of boron compounds on human reproduction. *Arch Toxicol*. 2020;94:717-24. doi: 10.1007/s00204-020-02700-x.
- ECHA (European Chemicals Agency). Annex XV dossier. Proposal for identification of a substance as substance of very high concern (SVHC). Substance name: Boric acid. EC Number: 233-139-2/234-343-4. CAS Number: 10043-35-3/11113-50-1. Helsinki: European Chemical Agency, 2010. <https://echa.europa.eu/de/registry-of-svhc-intentions/-/dislist/details/0b0236e180e4b39c>. Accessed 7 Jan 2020.
- Hasegawa R, Hirata-Koizumi M, Dourson ML, Parker A, Ono A, Hirose A. Safety assessment of boron by application of new uncertainty factors and their subdivision. *Regul Toxicol Pharmacol*. 2013;65:108–14. doi.org/10.1016/j.yrtph.2012.10.013.
- Krug AK, Kolde R, Gaspar JA, Rempel E, Balmer NV, Meganathan K, et al. Human embryonic stem cell-derived test systems for developmental neurotoxicity: a transcriptomics approach. *Arch Toxicol*. 2013;87:123-43. doi: 10.1007/s00204-012-0967-3.
- Leist M, Ghallab A, Graepel R, Marchan R, Hassan R, Bennekou SH, et al. Adverse outcome pathways: opportunities, limitations and open questions. *Arch Toxicol*. 2017;91:3477-3505. doi: 10.1007/s00204-017-2045-3.
- Moore JA. An assessment of boric acid and borax using the IEHR evaluative process for assessing human developmental and reproductive toxicity of agents. Expert scientific committee. *Reprod Toxicol*. 1997;11: 123–60. doi: 10.1016/S0890-6238(96)00204-3.
- Rempel E, Hoelting L, Waldmann T, Balmer NV, Schildknecht S, Grinberg M, Das Gaspar JA, et al. A transcriptome-based classifier to identify developmental toxicants by stem cell testing: design, validation and optimization for histone deacetylase inhibitors. *Arch Toxicol*. 2015;89:1599-618. doi: 10.1007/s00204-015-1573-y.
- Sachinidis A, Albrecht W, Nell P, Cherianidou A, Hewitt NJ, Edlund K, Hengstler JG. Road map for development of stem cell-based alternative test methods. *Trends Mol Med*. 2019;25:470-81. doi: 10.1016/j.molmed.2019.04.003.
- SCCS (Scientific Committee on Consumer Safety; EU Directorate for Health and Consumers). Opinion on boron compounds. Adopted 22 June 2010, Revision 28 Sep 2010. https://ec.europa.eu/health/scientific_committees/consumer_safety/docs/sccs_o_027.pdf. Accessed 7 Jan 2020.
- Shinde V, Stöber R, Nemade H, Sotiriadou I, Hescheler J, Hengstler J, et al. Transcriptomics of hepatocytes treated with toxicants for investigating molecular mechanisms underlying hepatotoxicity. *Methods Mol Biol*. 2015;1250:225-40. doi: 10.1007/978-1-4939-2074-7_16.
- Shinde V, Perumal Srinivasan S, Henry M, Rotshteyn T, Hescheler J, Rahnenführer J, et al. Comparison of a teratogenic transcriptome-based predictive test based on human embryonic versus inducible pluripotent stem cells. *Stem Cell Res Ther*. 2016;7:190. doi: 10.1186/s13287-016-0449-2.
- Shinde V, Hoelting L, Srinivasan SP, Meisig J, Megathan K, Jagtap S, et al. Definition of transcriptome-based indices for quantitative characterization of chemically disturbed stem cell development: introduction of the STOP-Tox_{unk} and STOP-Tox_{ukk} tests. *Arch Toxicol*. 2017;91:839-64. doi: 10.1007/s00204-016-1741-8.
- Waldmann T, Rempel E, Balmer NV, König A, Kolde R, Gaspar JA, et al. Design principles of concentration-dependent transcriptome deviations in drug-exposed differentiating stem cells. *Chem Res Toxicol*. 2014;27: 408-20. doi: 10.1021/tx400402j.
- Waldmann T, Grinberg M, König A, Rempel E, Schildknecht S, Henry M, et al. Stem cell transcriptome responses and corresponding biomarkers that indicate the transition from adaptive responses to cytotoxicity. *Chem Res Toxicol*. 2017;30:905-22. doi: 10.1021/acs.chemrestox.6b00259.
- Weng MK, Natarajan K, Scholz D, Ivanova VN, Sachinidis A, Hengstler JG, et al. Lineage-specific regulation of epigenetic modifier genes in human liver and brain. *PLoS One*. 2014;9:e102035. doi: 10.1371/journal.pone.0102035.